arXiv:1612.07500v1 [math-ph] 22 Dec 2016

ASYMPTOTICS OF DETERMINANTS OF 4-TH ORDER OPERATORS AT ZERO

ANDREY BADANIN AND EVGENY L. KOROTYAEV

Abstract. We consider fourth order ordinary differential operators with compactly supported coefficients on the half-line and on the line. The Fredholm determinant for this operator is an analytic function in the whole complex plane without zero. We describe the determinant at zero. We show that in the generic case it has a pole of order 4 in the case of the line and of order 1 in the case of the half-line.

1. Introduction and main results

We consider fourth order operators

$$H = H_0 + V, \quad \text{where} \quad V = 2\partial p \partial + q, \quad \partial = \frac{d}{dx},$$

and H_0 is one of the following:

Case 1: ∂^4 in $L^2(\mathbb{R}_+)$, with $y(0) = y''(0) = 0$ boundary conditions.

Case 2: ∂^4 in $L^2(\mathbb{R})$.

We assume that the functions p, q are compactly supported and belong to the space of functions $\mathcal{H}$, defined by

$$\mathcal{H} = \big\{ f \in L^1(\mathbb{R}_+) : \operatorname{supp} f \in [0, \gamma] \big\} \qquad \text{in Case 1}$$

$$\mathcal{H} = \big\{ \ f \in L^1(\mathbb{R}) \ : \operatorname{supp} f \in [0, \gamma] \big\} \qquad \text{in Case 2}$$

for some $\gamma > 0$. We consider the operator H_0 as unperturbed and the perturbed operator H is defined on the form domain $\mathrm{Dom}_{fd}(H) = \mathrm{Dom}_{fd}(H_0)$. The operator H has purely absolutely continuous spectrum $[0, \infty)$ plus a finite number of simple eigenvalues.

We rewrite V in the form

$$V = V_1 V_2, \qquad V_1 = (\partial |2p|^{\frac{1}{2}}, |q|^{\frac{1}{2}}), \qquad V_2 = \begin{pmatrix} (2p)^{\frac{1}{2}} \partial \\ q^{\frac{1}{2}} \end{pmatrix}, \tag{1.1}$$

and we set

$$R_0(k) = (H_0 - k^4)^{-1}, \qquad k \in \mathbb{K}_1 = \Big\{ k \in \mathbb{C} : \arg k \in \Big(0, \frac{\pi}{2}\Big) \Big\}. \tag{1.2}$$

Each operator $V_2 R_0(k) V_1, k \in \mathbb{K}_1$, is trace class and thus we can define a Fredholm determinant:

$$D(k) = \det \big(I + V_2 R_0(k) V_1 \big), \qquad k \in \mathbb{K}_1, \tag{1.3}$$

similar to second order operators. The function D is analytic in $\mathbb{K}_1$ and has an analytic extension from $\mathbb{K}_1$ into the whole complex plane without zero. Note that $k \in \overline{\mathbb{K}}_1 \setminus \{0\}$ is a zero of the determinant D iff $\lambda = k^4 \in \mathbb{R} \setminus \{0\}$ is an eigenvalue of the operator H. We define

Date: December 23, 2016.

1991 *Mathematics Subject Classification.* 47E05 (34L25).

Key words and phrases. fourth order operators, Fredholm determinant, asymptotics.

the resonances as zeros of the Fredholm determinant in $\mathbb{C} \setminus \overline{\mathbb{K}}_1$. A main goal of the present paper is to describe the determinant at the point zero.

Let $p, q \in \mathcal{H}$. Consider the equation

$$y'''' + 2(py')' + qy = k^4 y, \qquad k \in \mathbb{C}, \tag{1.4}$$

on the interval $[0, \gamma]$ in the class of function y satisfying the conditions

$$y^{[j]} \in AC([0,\gamma]), \quad j = 0, 1, 2, 3,$$

where $AC([0,\gamma])$ is the class of absolutely continuous functions on $[0,\gamma]$ and

$$y^{[0]} = y, \quad y^{[1]} = y', \quad y^{[2]} = y'', \quad y^{[3]} = y''' + 2py'.$$

Introduce the fundamental solutions $\varphi_j(x,k), j \in \mathbb{N}_4 = \{1,2,3,4\}$, of equation (1.4) satisfying the conditions

$$\varphi_j^{[\ell-1]}(0,k) = \delta_{j\ell}, \quad j, \ell \in \mathbb{N}_4. \tag{1.5}$$

Introduce the 4×4 matrix-valued function

$$M(k) = (M_{j\ell}(k))_{j,\ell=1}^4 = \big(\varphi_j^{[\ell-1]}(\gamma,k)\big)_{j,\ell=1}^4, \qquad k \in \mathbb{C}.$$

Each function $\varphi_j(x,\cdot), (j,x) \in \mathbb{N}_4 \times \mathbb{R}$, and M is entire in $\lambda = k^4$ and real at real λ.

Introduce the operators $H_j = \partial^4 + 2p\partial p + q, j = 1,2,3$, in $L^2(0,\gamma)$ with the boundary conditions

$$\begin{aligned} y(0) = y''(0) = y''(\gamma) = y^{[3]}(\gamma) = 0 \qquad &\text{for} \quad H_1 \\ y(0) = y''(0) = y'(\gamma) = y^{[3]}(\gamma) = 0 \qquad &\text{for} \quad H_2, \\ y''(0) = y^{[3]}(0) = y''(\gamma) = y^{[3]}(\gamma) = 0 \qquad &\text{for} \quad H_3, \end{aligned} \tag{1.6}$$

A spectrum $\sigma(H_j)$ of each operator $H_j, j = 1,2,3$ is discrete. Moreover,

$$\begin{aligned} \sigma(H_1) &= \{\lambda = k^4 \in \mathbb{C} : \Delta_{34}(k) = 0\}, \\ \sigma(H_2) &= \{\lambda = k^4 \in \mathbb{C} : \Delta_{24}(k) = 0\}, \\ \sigma(H_3) &= \{\lambda = k^4 \in \mathbb{C} : \Phi(k) = 0\}, \end{aligned} \tag{1.7}$$

where $\Phi(k), \Delta_{j\ell}(k), j, \ell \in \mathbb{N}_4$ are entire functions in $\lambda = k^4$, real at real λ, and given by

$$\Delta_{j\ell} = \det \begin{pmatrix} M_{j2} & M_{j4} \\ M_{\ell 2} & M_{\ell 4} \end{pmatrix}, \qquad \Phi = \det \begin{pmatrix} M_{31} & M_{32} \\ M_{41} & M_{42} \end{pmatrix}. \tag{1.8}$$

We prove the following main results.

Theorem 1.1. *Let $p, q \in \mathcal{H}$ in Case 1. Then the function $kD(k)$ is entire and satisfies:*

$$D(k) = \frac{e^{(i-1)k\gamma}}{2k}\Big((1+i)\Delta_{34}(0) + 2k\Delta_{24}(0) + (1-i)k^2\big(\Delta_{14}(0) + \Delta_{23}(0)\big) + O(k^3)\Big), \tag{1.9}$$

as $k \to 0$ uniformly on $\arg k \in [0, 2\pi]$. Moreover,

$$\big|\Delta_{34}(0)\big| + \big|\Delta_{24}(0)\big| + \big|\Delta_{14}(0) + \Delta_{23}(0)\big| \neq 0, \tag{1.10}$$

that is the function $k^{-2}D(k)$ has a pole of order $\geqslant 1$ at zero. Furthermore,

(i) The function $D(k)$ is entire iff $\lambda = 0$ is an eigenvalue of H_1,

(ii) The function $k^{-1}D(k)$ is entire iff $\lambda = 0$ is an eigenvalue of both H_1 and H_2.

Theorem 1.2. *Let $p, q \in \mathcal{H}$ in Case 2. Then the function $k^4 D(k)$ is entire and satisfies:*

$$D(k) = -\frac{1}{8k^4}\big(\Phi(0) + O(k)\big), \tag{1.11}$$

as $k \to 0$ uniformly on $\arg k \in [0, 2\pi]$. In particular, the function $k^3 D(k)$ is entire iff $\lambda = 0$ is an eigenvalue of H_3.

Remark. 1) It is well known that the Fredholm determinant for a Scrödinger operator on the half-line with compactly supported potential is an entire function, and the determinant may have a simple pole at zero in the case of a Scrödinger operator on the line.

2) Asymptotics (1.9) shows that the determinant D for the operator on the half-line has a pole of order $\leqslant 1$ at zero. Similarly, asymptotics (1.11) shows that the determinant D for the operator on the line has a pole of order $\leqslant 4$ at zero.

3) Coefficients of the asymptotics (1.9) are expressed in terms of auxiliary boundary problems. It gives a complete description of the determinant at zero in Case 1. The asymptotics (3.7) in Case 2 is more complicated. In this case we give necessary and sufficient conditions for the determinant to have a pole of maximal order 4 at zero.

4) The asymptotics (1.9), (1.11) were used for trace formulas in terms of resonances in the papers [BK16], [BK16x].

Jost [J47] expressed the determinant for a Schrödinger operator in terms of Wronskian of some specific solutions (Jost solutions). This relation is important to study spectral properties of the Schrödinger operator, inverse problems, and, in particular, resonances (zeros of the determinant). The resonances of Schrödinger operators studied by Zworski [Z87], Froese [F97], Hitrik [H99], Simon [S00], Korotyaev [K04], [K05].

Many papers are devoted to the scattering for one dimensional higher order $\geqslant 3$ operators, see the book [BDT88] and references therein. In connection with the Boussinesq equation the spectral problems for third order operators on the line considered by Deift, Tomei and Trubowitz [DTT82]. The corresponding periodic problem was studied by McKean [McK81]. Resonances for third order operators on the line were considered by Korotyaev [K16]. The spectral problems for fourth order operators on the line considered by Iwasaki [Iw88], [Iw88x], Hoppe, Laptev and Östensson [HLO06]. Resonances for fourth order operators considered by Badanin and Korotyaev [BK16], [BK16x]. Fourth order operators with periodic coefficients studied in [BK11].

The plan of the paper is as follows. In Section 2 we consider the operator on the half-line. The operator on the line is considered in Section 3.

2. Operator on the half-line

2.1. **Jost function.** We consider Case 1. Define the sets

$$\mathscr{S} = \Big\{k \in \mathbb{C} : \arg k \in \big(0, \frac{\pi}{4}\big)\Big\}, \qquad \mathscr{S}(r) = \Big\{k \in \mathscr{S} : |k| > r\Big\}, \quad r > 0. \tag{2.1}$$

Consider the equation on $\mathbb{R}_+$

$$y'''' + 2(py')' + qy = k^4 y, \qquad k \in \mathbb{C}, \tag{2.2}$$

in the class of function y satisfying the conditions

$$y^{[j]} \in AC_{loc}(\mathbb{R}_+), \quad j = 0, 1, 2, 3, \tag{2.3}$$

where $AC_{loc}(\mathbb{R}_+)$ is the class of locally absolutely continuous functions on $\mathbb{R}_+$. Let $\psi_1(x,k)$, $\psi_2(x,k)$, $(x,k)\in\mathbb{R}_+\times\mathscr{S}$, be Jost solutions of equation (2.2) satisfying the conditions

$$\psi_1(x,k)=e^{-kx},\quad \psi_2(x,k)=e^{ikx},\quad \text{as}\quad x>\gamma, \tag{2.4}$$

for all $k\in\mathscr{S}$. Each function $\psi_j(x,\cdot), j=1,2, x\in\mathbb{R}_+$, is analytic function of $k\in\mathscr{S}(r)$ for some $r>0$ large enough, see [Iw88]. Introduce the function $w(k)$ by

$$w(k)=\psi_1(0,k)\psi_2'(0,k)-\psi_1'(0,k)\psi_2(0,k),\qquad k\in\mathscr{S}(r). \tag{2.5}$$

Lemma 2.1. *Let $p,q\in\mathcal{H}$. Then the function $w(k)$ has an analytic extension from $\mathscr{S}(r)$ onto the whole complex plane and satisfies*

$$\begin{aligned} w(k)=-ke^{(i-1)k\gamma}\Big((1+i)\Delta_{34}(k)+2k\Delta_{24}(k)+(1-i)k^2\big(\Delta_{14}(k)+\Delta_{23}(k)\big)\\ -2ik^3\Delta_{13}(k)-(1+i)k^4\Delta_{12}(k)\Big),\quad \forall\quad k\in\mathbb{C}, \end{aligned} \tag{2.6}$$

where $\Delta_{j\ell}$ are given by (1.8). In particular, $w(0)=0$.

Proof. Let $k\in\mathscr{S}(r)$. We have the identities

$$\begin{pmatrix}\psi_j^{[0]}\\ \psi_j^{[1]}\\ \psi_j^{[2]}\\ \psi_j^{[3]}\end{pmatrix}(0)=M^{-1}\begin{pmatrix}\psi_j^{[0]}\\ \psi_j^{[1]}\\ \psi_j^{[2]}\\ \psi_j^{[3]}\end{pmatrix}(\gamma),\quad j=1,2, \tag{2.7}$$

here and below $\psi_j(x)=\psi_j(x,k),...$ The matrix M satisfies the identity

$$M^{-1}=-JM^{\top}J,\quad \text{where}\quad J=(J_{j\ell})_{j,\ell=1}^4=\begin{pmatrix}0&0&0&1\\0&0&-1&0\\0&1&0&0\\-1&0&0&0\end{pmatrix}.$$

Identities $J_{j\ell}=(-1)^{j+1}\delta_{j,5-\ell}$ yield

$$(M^{-1})_{jn}=-\sum_{\ell,m=1}^4 J_{j\ell}M_{m\ell}J_{mn}=-\sum_{\ell,m=1}^4(-1)^{j+m}\delta_{j,5-\ell}\delta_{m,5-n}M_{m\ell}=(-1)^{j+n}M_{5-n,5-j} \tag{2.8}$$

for all $j,n\in\mathbb{N}_4$. Identities (2.7), (2.8) give

$$\begin{aligned}\psi_j^{[m-1]}(0)=\sum_{n=1}^4(M^{-1})_{mn}\psi_j^{[n-1]}(\gamma)&=\sum_{n=1}^4(-1)^{m+n}M_{5-n,5-m}\psi_j^{[n-1]}(\gamma)\\ &=\sum_{\ell=1}^4(-1)^{m+\ell-1}M_{\ell,5-m}\psi_j^{[4-\ell]}(\gamma),\end{aligned} \tag{2.9}$$

for all $j=1,2, m\in\mathbb{N}_4$. Using the identities $\psi_1(x)=e^{-kx},\psi_2(x)=e^{ikx}$ as $x>\gamma$, and (2.3) we obtain

$$\psi_1^{[\ell-1]}(\gamma)=(-k)^{\ell-1}e^{-k\gamma},\qquad \psi_2^{[\ell-1]}(\gamma)=(ik)^{\ell-1}e^{ik\gamma},\qquad \ell\in\mathbb{N}_4.$$

Then for $m\in\mathbb{N}_4$ the identities (2.9) give

$$\psi_1^{[m-1]}(0)=(-1)^{m-1}e^{-k\gamma}\sum_{\ell=1}^4k^{4-\ell}M_{\ell,5-m},\quad \psi_2^{[m-1]}(0)=(-1)^{m-1}e^{ik\gamma}\sum_{\ell=1}^4(-ik)^{4-\ell}M_{\ell,5-m}.$$

Substituting these identities into (2.5) we obtain

$$w(k) = e^{(i-1)k\gamma} \det \begin{pmatrix} \sum\limits_{\ell=1}^{4} k^{4-\ell} M_{\ell 4}(k) & \sum\limits_{\ell=1}^{4} (-ik)^{4-\ell} M_{\ell 4}(k) \\ \sum\limits_{j=1}^{4} k^{4-j} M_{j2}(k) & \sum\limits_{j=1}^{4} (-ik)^{4-j} M_{j2}(k) \end{pmatrix}$$

$$= e^{(i-1)k\gamma} \sum_{j,\ell=1}^{4} (i^j - i^\ell) k^{8-j-\ell} M_{j2}(k) M_{\ell 4}(k) = e^{(i-1)k\gamma} \sum_{1 \leqslant j < \ell \leqslant 4} (i^j - i^\ell) k^{8-j-\ell} \Delta_{j\ell}(k),$$

which yields (2.6). ■

2.2. **Resolvents.** Equation (1.4) may be rewritten in the vector form

$$\mathbf{y}' = Q\mathbf{y}, \qquad \text{where} \quad \mathbf{y} = \begin{pmatrix} y^{[0]} \\ y^{[1]} \\ y^{[2]} \\ y^{[3]} \end{pmatrix}, \qquad Q = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2p & 0 & 1 \\ k^4 - q & 0 & 0 & 0 \end{pmatrix}. \tag{2.10}$$

Introduce the matrix-valued function

$$X(x,k) = \begin{pmatrix} \varphi_2^{[0]} & \varphi_4^{[0]} & \psi_1^{[0]} & \psi_2^{[0]} \\ \varphi_2^{[1]} & \varphi_4^{[1]} & \psi_1^{[1]} & \psi_2^{[1]} \\ \varphi_2^{[2]} & \varphi_4^{[2]} & \psi_1^{[2]} & \psi_2^{[2]} \\ \varphi_2^{[3]} & \varphi_4^{[3]} & \psi_1^{[3]} & \psi_2^{[3]} \end{pmatrix} (x,k), \qquad (x,k) \in \mathbb{R}_+ \times \mathscr{S}(r). \tag{2.11}$$

The function X satisfies equation (2.10). The conditions (1.5) and the definition (2.5) imply that the function $\det X(x,\cdot)$ has an analytic extension from $\mathscr{S}(r)$ onto the whole complex plane and satisfies

$$\det X(x,k) = \det X(0,k) = -w(k) \qquad \forall \quad (x,k) \in \mathbb{R}_+ \times \mathbb{C}. \tag{2.12}$$

Let $(x,k) \in \mathbb{R}_+ \times \mathscr{S}(r)$ for some $r > 0$ large enough. Then $\det X(x,k) \neq 0$ and we introduce the matrix-valued functions

$$\Gamma(x,k) = \big(\Gamma_{j\ell}(x,k)\big)_{j,\ell=1}^{4} = X^{-1}(x,k), \tag{2.13}$$

$$\Omega(x,k) = \big(\Omega_{j\ell}(x,k)\big)_{j,\ell=1}^{4} = \Gamma(x,k)\dot{X}(x,k) = X^{-1}(x,k)\dot{X}(x,k), \tag{2.14}$$

where $\dot{f} = \frac{\partial f}{\partial k}$.

Lemma 2.2. *Let $p, q \in \mathcal{H}, (x,k) \in \mathbb{R}_+ \times \mathscr{S}(r)$. Then the matrix-valued function $\Omega(x,k)$ satisfies*

$$\Omega' = 4k^3 \begin{pmatrix} \Gamma_{14}\varphi_2 & \Gamma_{14}\varphi_4 & \Gamma_{14}\psi_1 & \Gamma_{14}\psi_2 \\ \Gamma_{24}\varphi_2 & \Gamma_{24}\varphi_4 & \Gamma_{24}\psi_1 & \Gamma_{24}\psi_2 \\ \Gamma_{34}\varphi_2 & \Gamma_{34}\varphi_4 & \Gamma_{34}\psi_1 & \Gamma_{34}\psi_2 \\ \Gamma_{44}\varphi_2 & \Gamma_{44}\varphi_4 & \Gamma_{44}\psi_1 & \Gamma_{44}\psi_2 \end{pmatrix}. \tag{2.15}$$

Proof. Differentiating the equation $X' = QX$ with respect to k we obtain $\dot{X}' = Q\dot{X} + \dot{Q}X$. Moreover, the definition (2.13) gives

$$\Gamma' = (X^{-1})' = -X^{-1}X'X^{-1} = -X^{-1}Q = -\Gamma Q.$$

Then the definition (2.14) yields

$$\Omega' = (\Gamma\dot{X})' = \Gamma'\dot{X} + \Gamma\dot{X}' = -\Gamma Q\dot{X} + \Gamma(Q\dot{X} + \dot{Q}X) = \Gamma\dot{Q}X. \tag{2.16}$$

The definition (2.10) of Q gives

$$\dot{Q} = 4k^3 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Substituting this identity and (2.11) into the identity (2.16) we obtain (2.15). ■

We need the following result.

Lemma 2.3. *Let $p, q \in \mathcal{H}, k \in \mathscr{S}(r)$. Then*

i) The resolvent $(H - k^4)^{-1}$ is an integral operator with the kernel $R(x,t,k), x, t \in \mathbb{R}_+$, given by

$$R(x,t,k) = \begin{cases} -\Gamma_{14}(t,k)\varphi_2(x,k) - \Gamma_{24}(t,k)\varphi_4(x,k), & x < t \\ \Gamma_{34}(t,k)\psi_1(x,k) + \Gamma_{44}(t,k)\psi_2(x,k), & x > t \end{cases}. \tag{2.17}$$

ii) The kernel $R(x,t,k)$ satisfies

$$R(x,x,k) = -\frac{F_1'(x,k)}{4k^3} = \frac{F_2'(x,k)}{4k^3} \tag{2.18}$$

for all $x \in \mathbb{R}_+$, where

$$F_1 = \Omega_{11} + \Omega_{22}, \qquad F_2 = \Omega_{33} + \Omega_{44}. \tag{2.19}$$

Moreover,

$$F_1(0,k) = 0, \qquad F_2(x,k) = F_2^0(x,k) \quad \text{as} \quad x > \gamma, \tag{2.20}$$

where F_2^0 is equal to F_2 for $p = q = 0$.

Proof. i) Let $(t,k) \in \mathbb{R}_+ \times \mathscr{S}$. The function $G(x) = R(x,t,k)$ satisfies

$$G'''' + 2(pG')' + qG = k^4 G, \qquad \forall \quad x \in \mathbb{R}_+ \setminus \{t\},$$
$$G, G', G'' \in AC_{loc}(\mathbb{R}_+), \qquad G^{[3]} \in AC_{loc}(0,t) \cap AC_{loc}(t, +\infty),$$
$$G^{[3]}(t+0) - G^{[3]}(t-0) = 1, \qquad G(0) = G''(0) = 0, \qquad \lim_{x\to\infty} G(x) = 0.$$

These conditions give

$$R(x,t,k) = \begin{cases} -A_1(t,k)\varphi_2(x,k) - A_2(t,k)\varphi_4(x,k), & x < t \\ A_3(t,k)\psi_1(x,k) + A_4(t,k)\psi_2(x,k), & x > t \end{cases},$$

where $A_j(t,k), j \in \mathbb{N}_4$ satisfy

$$X(t,k) \begin{pmatrix} A_1 \\ A_2 \\ A_3 \\ A_4 \end{pmatrix} (t,k) = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$

The definition (2.13) gives $A_j = \Gamma_{j4}, j \in \mathbb{N}_4$, which yields (2.17).

ii) The identities (2.15) and (2.17) imply (2.18).

Let $x > \gamma$. Then $p = q = 0$ and $\Omega(x,k) = \Omega^0(x,k)$, where $\Omega^0(x,k)$ is equal to $\Omega(x,k)$ for $p = q = 0$. This yields the second identity in (2.20).

The identity (2.11) gives $\dot X_{\ell j}(0,k) = 0, \ell \in \mathbb{N}_4, j = 1,2$. The definitions (2.14), (2.19) imply $F_1 = (\Gamma \dot X)_{11} + (\Gamma \dot X)_{22}$. Then

$$F_1(0,k) = \sum_{\ell=1}^{4} \big(\Gamma_{1\ell}(0,k)\dot X_{\ell 1}(0,k) + \Gamma_{2\ell}(0,k)\dot X_{\ell 2}(0,k)\big) = 0,$$

which yields the first identity in (2.20). ■

2.3. **Determinant.** Now we express the determinant $D(k)$ in terms of the Jost function $w(k)$.

Lemma 2.4. *Let $p, q \in \mathcal{H}$. Then the determinant D, given by (1.3), satisfies*

$$D(k) = -\frac{w(k)}{2k^2} \qquad \forall \quad k \in \mathbb{C} \setminus \{0\}, \tag{2.21}$$

where w is given by (2.5). Moreover,

$$D = \frac{e^{(i-1)k\gamma}}{2k}\Big((1+i)\Delta_{34} + 2k\Delta_{24} + (1-i)k^2\big(\Delta_{14}+\Delta_{23}\big) - 2ik^3\Delta_{13} - (1+i)k^4\Delta_{12}\Big) \tag{2.22}$$

on $\mathbb{C} \setminus \{0\}$, where $\Delta_{j\ell}$ are given by (1.8).

Proof. Let $0 < x < \gamma < \alpha < \infty$ and let $k \in \mathscr{S}(r)$. Integrating the first identity in (2.18) in the interval $(0,x)$ and the second identity in (2.18) in the interval (x,α) and using (2.19) and (2.20) we obtain

$$\int_0^\alpha R(t,t,k)dt = \frac{1}{4k^3}\Big(\int_x^\alpha F_2'(t,k)dt - \int_0^x F_1'(t,k)dt\Big) = -\frac{1}{4k^3}\big(\operatorname{Tr}\Omega(x,k) - F_2^0(\alpha,k)\big). \tag{2.23}$$

Furthermore, identities (2.12) and (2.13) imply

$$\operatorname{Tr}\Omega = \operatorname{Tr}\Gamma\dot X = \operatorname{Tr} X^{-1}\dot X = \frac{1}{\det X}\frac{\partial \det X}{\partial k} = \frac{\dot w}{w}. \tag{2.24}$$

Substituting (2.24) into (2.23) we obtain

$$\int_0^\alpha R(t,t,k)dt = -\frac{1}{4k^3}\Big(\frac{\dot w(k)}{w(k)} - F_2^0(\alpha,k)\Big),$$

Using a similar identity for $p = q = 0$ we obtain

$$\operatorname{Tr}\big(R(k) - R_0(k)\big) = \int_0^\alpha \big(R(t,t,k) - R_0(t,t,k)\big)dt = -\frac{1}{4k^3}\Big(\frac{\dot w(k)}{w(k)} - \frac{\dot w_0(k)}{w_0(k)}\Big),$$

where R_0, w_0 are equal to R, w as $p = q = 0$ respectively. The identity

$$\frac{D'(k)}{D(k)} = \operatorname{Tr} R_0(k) V R(k) = \operatorname{Tr}(R_0(k) - R(k)),$$

yields

$$D(k) = \frac{w(k)}{w_0(k)}.$$

The identity $w_0(k) = -2k^2$ gives (2.21). The identities (2.6) and (2.21) imply (2.22). ■

We are ready to prove Theorem 1.1.

Proof of Theorem 1.1. The identity (2.22) yields the asymptotics (1.9).

We prove (1.10). Assume that $\Delta_{34}(0) = \Delta_{24}(0) = \Delta_{14}(0) + \Delta_{23}(0) = 0$. Then the definitions (1.8) give

$$\det \begin{pmatrix} M_{32} & M_{34} \\ M_{42} & M_{44} \end{pmatrix}(0) = \det \begin{pmatrix} M_{22} & M_{24} \\ M_{42} & M_{44} \end{pmatrix}(0) = \det \begin{pmatrix} M_{12} & M_{14} \\ M_{42} & M_{44} \end{pmatrix}(0) + \det \begin{pmatrix} M_{22} & M_{24} \\ M_{32} & M_{34} \end{pmatrix}(0) = 0.$$

These identities imply

$$\det \begin{pmatrix} M_{22} & M_{24} \\ M_{32} & M_{34} \end{pmatrix}(0) = \det \begin{pmatrix} M_{12} & M_{14} \\ M_{42} & M_{44} \end{pmatrix}(0) = 0.$$

Then $\det M(0) = 0$. It is in contradiction with the Liouville identity $\det M = 1$, which proves (1.10).

(i) The asymptotics (1.9) gives that the function D is entire iff $\Delta_{34}(0) = 0$. The first identity in (1.7) shows that in this case $\lambda = 0$ is an eigenvalue of the operator H_1.

(ii) The asymptotics (1.9) gives that the function $k^{-1}D$ is entire iff $\Delta_{34}(0) = \Delta_{24}(0) = 0$. The first and second identities in (1.7) show that in this case $\lambda = 0$ is an eigenvalue of the operators H_1 and H_2. ■

2.4. **Square of a second order operator.** Now we consider the square of a second order operator.

Proposition 2.5. *Let $p, p'' \in L^1(\mathbb{R}_+)$ and $\operatorname{supp} p \in [0, \gamma]$. Let $H = h^2$, where h is the operator $h = -\partial^2 - p$ on $\mathbb{R}_+$ with the condition $y(0) = 0$. Then the determinant D satisfies the identity*

$$D(k) = \psi(0, ik)\psi(0, k) \qquad \forall \quad k \in \mathbb{C}, \tag{2.25}$$

where $\psi(x, k), (x, k) \in \mathbb{R}_+ \times \mathbb{C}$, is the solution of the equation $-y'' - py = zy$, satisfying the identity

$$\begin{gathered} \psi(x, k) = e^{ikx} \qquad \forall \quad x > \gamma, \\ k = \sqrt{z}, \quad k \in \mathbb{C}_+ \quad \text{as} \quad z \in \mathbb{C} \setminus \mathbb{R}_+. \end{gathered} \tag{2.26}$$

Proof. In the considered case we have $\psi_1(x, k) = \psi(x, ik), \psi_2(x, k) = \psi(x, k)$. Using the identities $\psi''(x, k) = (-p(x) - k^2)\psi(x, k)$ we obtain

$$\begin{gathered} w(k) = \psi_1(0, k)\psi_2'(0, k) - \psi_1'(0, k)\psi_2(0, k) = \psi(0, ik)\psi''(0, k) - \psi(0, k)\psi''(0, ik) \\ = \psi(0, ik)(-p(0) - k^2)\psi(0, k) - \psi(0, k)(-p(0) + k^2)\psi(0, ik) = -2k^2\psi(0, ik)\psi(0, k). \end{gathered}$$

The identity (2.21) yields (2.25). ■

Remark. The identity (2.25) used in [BK16] in order to describe resonances of h^2.

3. Operator on the line

3.1. **Jost function.** We consider Case 2. Let $\mathscr{S}, \mathscr{S}(r), r > 0$ be the sets given by the relations (2.1). Consider the equation on $\mathbb{R}$

$$y'''' + 2(py')' + qy = k^4 y, \qquad k \in \mathbb{C}, \tag{3.1}$$

in the class of functions y satisfying the conditions

$$y^{[j]} \in AC_{loc}(\mathbb{R}), \quad j = 0, 1, 2, 3.$$

Let $\psi_j(x,k), j\in\mathbb{N}_4$, be Jost solutions of equation (3.1) satisfying the conditions

$$\begin{aligned}\psi_1(x,k)&=e^{-kx},\quad \psi_2(x,k)=e^{ikx},\quad \text{as}\quad x>\gamma,\\ \psi_3(x,k)&=e^{-ikx},\quad \psi_4(x,k)=e^{kx},\quad \text{as}\quad x<0,\end{aligned}\tag{3.2}$$

for all $k\in\mathscr{S}$. Each function $\psi_j(x,\cdot),(j,x)\in\mathbb{N}_4\times\mathbb{R}$, is analytic in $k\in\mathscr{S}(r)$ for some $r>0$ large enough, see [Iw88].

Let $k\in\mathscr{S}(r)$. Introduce the Wronskian $w(k)$ by

$$w(k)=\det\begin{pmatrix}\psi_1^{[0]}&\psi_2^{[0]}&\psi_3^{[0]}&\psi_4^{[0]}\\ \psi_1^{[1]}&\psi_2^{[1]}&\psi_3^{[1]}&\psi_4^{[1]}\\ \psi_1^{[2]}&\psi_2^{[2]}&\psi_3^{[2]}&\psi_4^{[2]}\\ \psi_1^{[3]}&\psi_2^{[3]}&\psi_3^{[3]}&\psi_4^{[3]}\end{pmatrix}(x,k),\quad k\in\mathscr{S}(r).\tag{3.3}$$

Östensson and Yafaev [OY12] proved that the function w has an analytic extension from $\mathscr{S}(r)$ onto the whole complex plane and

$$D(k)=-\frac{w(k)}{16ik^6}\qquad\forall\quad k\in\mathbb{C}\setminus\{0\}.\tag{3.4}$$

This formula shows that the determinant may have a pole of order $\leqslant 6$ at zero. We prove that the order is $\leqslant 4$.

Introduce the functions $\Delta_{jn,\ell m}(k), 1\leqslant j<n\leqslant 4, 1\leqslant\ell<m\leqslant 4, k\in\mathbb{C}$ by

$$\Delta_{jn,\ell m}=\det\begin{pmatrix}M_{j\ell}&M_{jm}\\ M_{n\ell}&M_{nm}\end{pmatrix}.$$

Each function $\Delta_{jn,\ell m}$ is an entire function of the variable $\lambda=k^4$.

Lemma 3.1. *Let $p,q\in\mathcal{H}$. Then the determinant $D(k)$ satisfies*

$$D(k)=-\frac{e^{(i-1)k\gamma}}{8k^4}\sum_{n=0}^{8}A_n(k)k^n,\qquad k\in\mathbb{C}\setminus\{0\},\tag{3.5}$$

where $A_n(k)$ are entire in $\lambda=k^4$ functions given by

$$\begin{aligned}A_0&=\Delta_{34,12},\qquad A_1=(1-i)(\Delta_{34,13}-\Delta_{24,12}),\\ A_2&=i(2\Delta_{24,13}-\Delta_{23,12}-\Delta_{14,12}-\Delta_{34,14}-\Delta_{34,23}),\\ A_3&=(1+i)(\Delta_{13,12}-\Delta_{23,13}-\Delta_{14,13}+\Delta_{24,14}+\Delta_{24,23}-\Delta_{34,24}),\\ A_4&=2\Delta_{24,24}-\Delta_{12,12}+2\Delta_{13,13}-\Delta_{23,14}-\Delta_{14,14}-\Delta_{23,23}-\Delta_{14,23}-\Delta_{34,34},\\ A_5&=(1-i)(\Delta_{24,34}-\Delta_{12,13}+\Delta_{13,14}+\Delta_{13,23}-\Delta_{23,24}-\Delta_{14,24}),\\ A_6&=i(\Delta_{12,14}+\Delta_{12,23}+\Delta_{23,34}+\Delta_{14,34}-2\Delta_{13,24}),\\ A_7&=(1+i)(\Delta_{12,24}-\Delta_{13,34}),\qquad A_8=\Delta_{12,34}.\end{aligned}\tag{3.6}$$

Moreover,

$$D(k)=-\frac{e^{(i-1)k\gamma}}{8k^4}\Big(A_0(0)+A_1(0)k+A_2(0)k^2+A_3(0)k^3+O(k^4)\Big),\tag{3.7}$$

as $k\to 0$ uniformly in $\arg k\in[0,2\pi]$.

Proof. Substituting the identities

$$\begin{pmatrix} \psi_j^{[0]} \\ \psi_j^{[1]} \\ \psi_j^{[2]} \\ \psi_j^{[3]} \end{pmatrix}(\gamma,k) = M(k) \begin{pmatrix} \psi_j^{[0]} \\ \psi_j^{[1]} \\ \psi_j^{[2]} \\ \psi_j^{[3]} \end{pmatrix}(0,k), \qquad j=3,4,$$

into the definition (3.3) we obtain

$$w(k) = \sum_{\ell,m=1}^{4} W_{\ell m}(k)\psi_3^{[\ell-1]}(0,k)\psi_4^{[m-1]}(0,k) = \sum_{1\leqslant \ell<m\leqslant 4} W_{\ell m}(k)E_{\ell m}(k), \tag{3.8}$$

where

$$E_{\ell m}(k) = \det \begin{pmatrix} \psi_3^{[\ell-1]} & \psi_3^{[m-1]} \\ \psi_4^{[\ell-1]} & \psi_4^{[m-1]} \end{pmatrix}(0,k), \tag{3.9}$$

$$W_{\ell m}(k) = \det \begin{pmatrix} \psi_1^{[0]}(\gamma,k) & \psi_2^{[0]}(\gamma,k) & M_{1\ell}(k) & M_{1m}(k) \\ \psi_1^{[1]}(\gamma,k) & \psi_2^{[1]}(\gamma,k) & M_{2\ell}(k) & M_{2m}(k) \\ \psi_1^{[2]}(\gamma,k) & \psi_2^{[2]}(\gamma,k) & M_{3\ell}(k) & M_{3m}(k) \\ \psi_1^{[3]}(\gamma,k) & \psi_2^{[3]}(\gamma,k) & M_{4\ell}(k) & M_{4m}(k) \end{pmatrix}. \tag{3.10}$$

Recall that $\psi_j^{[\ell-1]}(\cdot,k) \in AC_{loc}(\mathbb{R})$ for all $j,\ell \in \mathbb{N}_4$. Then the definitions (3.2) imply

$$\psi_1^{[j]}(\gamma,k) = (-k)^j e^{-k\gamma}, \quad \psi_2^{[j]}(\gamma,k) = (ik)^j e^{ik\gamma}, \quad \psi_3^{[j]}(0,k) = (-ik)^j, \quad \psi_4^{[j]}(0,k) = k^j, \tag{3.11}$$

for all $j \in \mathbb{N}_4$. Substituting the identities (3.11) into (3.9) we have

$$E_{\ell m} = \big((-i)^{\ell-1} - (-i)^{m-1}\big)k^{\ell+m-2},$$

The identity (3.8) gives

$$w = (1+i)kW_{12} + 2k^2W_{13} + (1-i)k^3\big(W_{14}+W_{23}\big) - 2ik^4W_{24} - (1+i)k^5W_{34}. \tag{3.12}$$

Substituting the identities (3.11) into (3.10) we obtain

$$W_{\ell m} = e^{(i-1)k\gamma} \det \begin{pmatrix} 1 & 1 & M_{1\ell} & M_{1m} \\ -k & ik & M_{2\ell} & M_{2m} \\ k^2 & -k^2 & M_{3\ell} & M_{3m} \\ -k^3 & -ik^3 & M_{4\ell} & M_{4m} \end{pmatrix} = e^{(i-1)k\gamma}\Big((1+i)k\Delta_{34,\ell m} - 2k^2\Delta_{24,\ell m}$$
$$+(1-i)k^3(\Delta_{23,\ell m}+\Delta_{14,\ell m}) + 2ik^4\Delta_{13,\ell m} - (1+i)k^5\Delta_{12,\ell m}\Big).$$

Substituting this identity into (3.12) we obtain

$$w = 2ik^2 e^{(i-1)k\gamma} \sum_{n=0}^{8} A_n k^n,$$

where A_n are given by (3.6). The identity (3.4) gives (3.5), which yields the asymptotics (3.7). ■

Proof of Theorem 1.2. The asymptotics (3.7) gives (1.11). The asymptotics (1.11) implies that the function D has a pole of order $\leqslant 3$ at $k=0$ iff $\Phi(0)=0$. The identity (1.7) shows that this is so iff $\lambda = 0$ is an eigenvalue of the operator H_3. ■

3.2. **Square of a second order operator.** Now we consider the square of a second order operator.

Proposition 3.2. *Let $p, p'' \in L^1(\mathbb{R})$ and $\operatorname{supp} p \in [0,\gamma]$. Let $H = h^2$, where h is the operator $h = -\partial^2 - p$ on $\mathbb{R}$. Then the determinant D satisfies the identity*

$$D(k) = \frac{i}{4k^2}\{\psi_+(x,k),\psi_-(x,k)\}\{\psi_+(x,ik),\psi_-(x,ik)\} \qquad \forall \quad k \in \mathbb{C}\setminus\{0\}, \tag{3.13}$$

where $\{f,g\} = fg' - f'g$ and $\psi_\pm(x,k), (x,k) \in \mathbb{R}\times\mathbb{C}$, are solutions of the equation $-y'' - py = zy$, satisfying the identity

$$\begin{gathered}\psi_+(x,k) = e^{ikx} \quad \forall \quad x > \gamma, \qquad \psi_-(x,k) = e^{-ikx} \quad \forall \quad x < 0,\\ k = \sqrt{z}, \quad k \in \mathbb{C}_+ \quad \text{as} \quad z \in \mathbb{C}\setminus\mathbb{R}_+.\end{gathered} \tag{3.14}$$

Proof. In the considered case we have

$$\psi_1(x,k) = \psi_+(x,ik), \quad \psi_2(x,k) = \psi_+(x,k), \quad \psi_3(x,k) = \psi_-(x,k), \quad \psi_4(x,k) = \psi_-(x,ik).$$

Substituting the identities $\psi''_\pm(x,k) = (-p(x) - k^2)\psi_\pm(x,k)$ into (3.3) we obtain

$$w(k) = k^4 \det \begin{pmatrix} \psi_+(x,ik) & \psi_+(x,k) & \psi_-(x,k) & \psi_-(x,ik) \\ \psi'_+(x,ik) & \psi'_+(x,k) & \psi'_-(x,k) & \psi'_-(x,ik) \\ \psi_+(x,ik) & -\psi_+(x,k) & -\psi_-(x,k) & \psi_-(x,ik) \\ \psi'_+(x,ik) & -\psi'_+(x,k) & -\psi'_-(x,k) & \psi'_-(x,ik) \end{pmatrix},$$

which yields

$$w(k) = 4k^4\{\psi_+(x,k),\psi_-(x,k)\}\{\psi_+(x,ik),\psi_-(x,ik)\}.$$

The identity (3.4) gives (3.13). ■

Remark. The identity (3.13) used in [BK16x] in order to describe resonances of h^2.

Acknowledgments. The study was supported by the RNF grant No 15-11-30007.

References

[BK16] Badanin, A., Korotyaev, E. Resonances for Euler-Bernoulli operator on the half-line. Preprint, 2016.

[BK16x] Badanin, A., Korotyaev, E. Resonances of 4-th Order Differential Operators. Preprint, 2016.

[BK11] Badanin, A., Korotyaev, E. Spectral estimates for periodic fourth order operators. St.Petersburg Math. J. 22:5 (2011) 703–736.

[BDT88] Beals, R., Deift, P., Tomei, C. Direct and inverse scattering on the line, Mathematical survays and monograph series, No. 28, AMS, Providence, 1988.

[DTT82] Deift, P., Tomei, C., Trubowitz, E. Inverse scattering and the Boussinesq equation, Comm. on Pure and Appl. Math. 35 (1982) 567–628.

[F97] Froese, R. Asymptotic distribution of resonances in one dimension. Journal of differential equations, 137 (1997), 251–272.

[H99] Hitrik, M. Bounds on scattering poles in one dimension. Comm. Math. Phys. 208 (1999), no. 2, 381–411.

[HLO06] Hoppe, J., Laptev, A., Östensson, J. Solitons and the removal of eigenvalues for fourth-order differential operators, Int. Math. Res. Not., (2006), 14 pp.

[Iw88] Iwasaki, K. Scattering Theory for 4-th Order Differential Operators, I. Japanese journal of mathematics. New series, 14(1) (1988), 1–57.

[Iw88x] Iwasaki, K. Scattering theory for 4-th order differential operator, II. Japanese journal of mathematics. New series, 14(1) (1988), 59–96.

[J47] Jost, R. Uber die falschen Nullstellen der Eigenwerte der S-Matrix. Helvetica Physica Acta, 20(3) (1947), 256–266.

[K04] Korotyaev, E. Inverse resonance scattering on the half line. Asymptot. Anal. 37 (2004), no. 3-4, 215–226.

[K05] Korotyaev, E. Inverse resonance scattering on the real line. Inverse Problems 21.1 (2005), 325–341.
[K16] Korotyaev, E. Resonances of third order differential operators. ArXiv preprint arXiv:1605.01842 (2016).
[McK81] McKean, H. Boussinesq's equation on the circle, Com. Pure and Appl. Math. 34 (1981) 599–691.
[OY12] Östensson, J., Yafaev, D. R. A trace formula for differential operators of arbitrary order. In A panorama of modern operator theory and related topics. Springer Basel, 2012, 541–570.
[S00] Simon, B. Resonances in one dimension and Fredholm determinants, J. Funct. Anal. 178 (2000), no. 2, 396–420.
[Z87] Zworski, M. Distribution of poles for scattering on the real line, J. Funct. Anal. 73(1987), 277–296.

Saint-Petersburg State University, Universitetskaya nab. 7/9, St. Petersburg, 199034 Russia, an.badanin@gmail.com, a.badanin@spbu.ru, korotyaev@gmail.com, e.korotyaev@spbu.ru